

AD-A042 865

AEROSPACE CORP EL SEGUNDO CALIF SPACE SCIENCES LAB

F/G 4/1

K-SHELL X-RAY FLUXES AT SATELLITE ALTITUDE FROM PROTON PRECIPIT--ETC(U)

JUL 77 J G LUHMANN , J B BLAKE

F04701-76-C-0077

UNCLASSIFIED

TR-0077(2260-20)-12


SAMSO-TR-77-140

NL

| OF |  
ADA042865



END  
DATE  
FILMED  
9-77  
DDC



AD A 042865

## K-Shell X-Ray Fluxes at Satellite Altitude From Proton Precipitation

Space Sciences Laboratory  
The Ivan A. Getting Laboratories  
The Aerospace Corporation  
El Segundo, Calif. 90245


15 July 1977

Interim Report

APPROVED FOR PUBLIC RELEASE:  
DISTRIBUTION UNLIMITED

Prepared for  
SPACE AND MISSILE SYSTEMS ORGANIZATION  
AIR FORCE SYSTEMS COMMAND  
Los Angeles Air Force Station  
P.O. Box 92960, Worldway Postal Center  
Los Angeles, Calif. 90009

AD NO. \_\_\_\_\_  
DDC FILE COPY



This interim report was submitted by The Aerospace Corporation, El Segundo, CA 90245, under Contract No. F04701-76-C-0077 with the Space and Missile Systems Organization, Deputy for Advanced Space Programs, P.O. Box 92960, Worldway Postal Center, Los Angeles, CA 90009. It was reviewed and approved for The Aerospace Corporation by G. A. Paulikas, Director, Space Sciences Laboratory. Lieutenant Dara Batki, SAMSO/YAPT, was the project officer for Advanced Space Programs.

This report has been reviewed by the Information Office (OI) and is releasable to the National Technical Information Service (NTIS). At NTIS, it will be available to the general public, including foreign nations.

This technical report has been reviewed and is approved for publication. Publication of this report does not constitute Air Force approval of the report's findings or conclusions. It is published only for the exchange and stimulation of ideas.

Dara Batki  
Dara Batki, 2nd Lt, USAF  
Project Officer

Robert W. Lindemuth  
Robert W. Lindemuth, Lt Col, USAF

FOR THE COMMANDER

Floyd R. Stuart  
Floyd R. Stuart, Colonel, USAF  
Deputy for Advanced Space Programs

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

19 REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM	
1. REPORT NUMBER	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER	
18 SAMSQ-TR-77-140		9	
4. TITLE (and Subtitle)	5. TYPE OF REPORT & PERIOD COVERED		
6 K-SHELL X-RAY FLUXES AT SATELLITE ALTITUDE FROM PROTON PRECIPITATION	Interim <i>rept.</i>		
7. AUTHOR(s)	6. PERFORMING ORG. REPORT NUMBER		
10 Janet G. Luhmann and J. Bernard Blake	14 TR-0077(2260-20)-12		
9. PERFORMING ORGANIZATION NAME AND ADDRESS		8. CONTRACT OR GRANT NUMBER(s)	
The Aerospace Corporation El Segundo, Calif. 90245		15 F04701-76-C-0077	
11. CONTROLLING OFFICE NAME AND ADDRESS		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS	
Space and Missile Systems Organization Air Force Systems Command Los Angeles, Calif. 90009			
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE	
12 12p.		11 15 July 1977	
		13. NUMBER OF PAGES	
		10	
		15. SECURITY CLASS. (of this report)	
		Unclassified	
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this Report)			
Approved for public release; distribution unlimited			
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)			
AUG 15 1977			
18. SUPPLEMENTARY NOTES			
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)			
Proton Precipitation X rays			
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)			
A sample calculation is presented of K-shell x-ray emission resulting from energetic proton precipitation into the upper atmosphere. It is found that intense fluxes of K-shell x-rays at low satellite altitudes can result from observed fluxes of precipitating protons.			

DD FORM 1473  
(FACSIMILE)

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

407512

JB



## CONTENTS

1.	INTRODUCTION . . . . .	3
2.	CALCULATIONS . . . . .	5
3.	DISCUSSION . . . . .	9
	REFERENCES . . . . .	11

## FIGURES

1.	K-Shell X-Ray Cross Sections . . . . .	6
2.	Satellite Measurements of Proton and Electron Fluxes on 14 April 1969 . . . . .	7
3.	Proton Energy Spectrum as observed at 75570 sec UT on 14 April 1969 . . . . .	7

ACCESSION NO.	
NTIS	<input checked="" type="checkbox"/> Section 1
DDC	<input type="checkbox"/> Section 2
UNANNOUNCED	<input type="checkbox"/>
JUL 1 1969	
BY	
DISSEMINATION CODES	
Dr	SPECIAL
A.	

## 1. INTRODUCTION

Luhmann and Blake (1977) have presented sample calculations of the soft bremsstrahlung and K-shell x-rays (0.10 keV to 10 keV) present at low satellite altitude resulting from the precipitation of electrons into the upper atmosphere. Since proton precipitation occurs also, it is of interest to determine if such precipitation results in a detectable x-ray signal. It happens that the proton bremsstrahlung cross-section is negligible but the cross-section for producing K-shell x-rays by proton impact upon nitrogen and oxygen is significant above a few tens of keV.

In this paper a sample calculation is presented based upon a satellite measurement of proton precipitation. The general question of the detectability of x-rays from proton precipitation is discussed briefly.

## 2. CALCULATIONS

The cross-section for the production of K-shell x-rays by protons impinging upon oxygen, nitrogen and argon are required for the calculation. Measured cross sections have been presented by Hart et al. (1969) for protons with energies between 20 keV and 100 keV impinging on oxygen. Data are available also for the production of K-shell x-rays by protons with energies between 15 keV and 1900 keV impacting carbon, magnesium and other heavier elements (Khan et al., 1965). In the absence of measurements, these latter data were used in concert with well-understood theoretical principles (Merzbacher and Lewis, 1958; Madison and Merzbacher, 1975) to provide the cross-sections for K-shell x-ray production by protons with energies from 20 keV to 700 keV impinging upon nitrogen, oxygen and argon.

The calculation was performed as described in Luhmann and Blake (1977) by substituting the proton K-shell cross-sections described above for the electron cross-sections given by Tawara et al. (1973), replacing an electron range-energy relationship by one for protons, and deleting the bremsstrahlung formalism.

The cross-sections used in the calculation are shown in Figure 1. It is clear from Figure 1 that proton energies of several tens of keV are required to excite appreciable K-shell emission in oxygen and nitrogen and that hundreds of keV are required in the case of argon.






Fig. 1. K-Shell X-Ray Cross-Sections (p,x).

The proton precipitation flux, spectra and pitch-angle distribution used in this sample calculation were observed during the solar particle event which began on 11 April 1969; the specific data were obtained at 75570 sec UT on 14 April. These measurements are described in detail by Blake and Vampola (1974). Figures 2 and 3 are adapted from their paper. Figure 2 shows the measured proton and electron fluxes during a polar cap transverse by the OV1-19 satellite. The proton energy spectrum observed at 75570 sec UT is shown in Figure 3. Note that at the location of the peak proton precipitation intensities, the energetic electron ( $E > 100$  keV) precipitation is nil. Unfortunately, no measurements of auroral energy electrons were made aboard the spacecraft, and thus possible K-shell x-ray emission due to auroral electron precipitation cannot be calculated.




Fig. 2. Satellite Measurements of Proton and Electron Fluxes on 14 April 1969.




Fig. 3. Proton Energy Spectrum as observed at 75570 sec UT on 14 April 1969. (See Fig. 2.)

The present calculation yields the following K-shell x-ray fluxes at satellite altitude due to the observed proton spectrum shown in Figure 3:

Nitrogen:	$4 \times 10^7 \text{ cm}^2\text{-sec;}$
Oxygen:	$2 \times 10^7 / \text{cm}^2\text{-sec;}$
Argon:	$7 \times 10^{-2} / \text{cm}^2\text{-sec.}$

The absence of significant argon K-shell x-rays is due to the fact that the proton-argon cross-section does not become appreciable (Figure 1) until an energy where there was little proton precipitation during the event depicted in Figure 2.

### 3. DISCUSSION

It is clear from the energy dependence of the K-shell cross-sections that normal auroral-energy proton precipitation will not cause significant K-shell x-ray emission from upper atmospheric nitrogen or oxygen. However, the sample calculation does show that intense fluxes of K-shell x-rays from nitrogen and oxygen result from the more energetic proton precipitation events that have been observed. (It should be noted that the proton precipitation shown in Figure 2, although occurring during a solar-particle event, does not simply consist of solar protons. The protons were accelerated by magnetospheric processes. Whether or not they were solar protons, with energies well above typical solar wind energies before magnetospheric acceleration, could not be determined (Blake and Vampola, 1974)).

Since the protons generate no bremsstrahlung continuum, x-ray emission from precipitating protons and that from precipitating electrons differs in a readily detectable fashion. If only K-shell lines are present, the precipitation consists of protons, not electrons. The shape of the bremsstrahlung spectrum, if present, reveals some information about the spectrum of the precipitating energetic electrons (Imhof et al., 1975a, 1975b; Luhmann and Blake, 1977). Thus satellite x-ray measurements which span the energy range from  $\sim 300$  eV to  $\sim 300$  keV will permit remote sensing of electron and energetic proton precipitation.



#### 4. REFERENCES

- |  |       |   |
|--|-------|---|
| Blake, J. B. and Vampola, A. L.                            | 1974  | in Correlated interplanetary and Magnetospheric Observations, (ed. D.E. Page), 173, Reidel, Dorchecht, Holland. |
| Hart, R. R., Reuter, F. W. and Smith, H. P.                | 1969  | Phys. Rev. <u>179</u> , 4.  |
| Imhof, W. L., Nakano, G. H. Gains, E. E. and Reagan, J. B. | 1975a | J. Geophys. Res. <u>80</u> , 3622.  |
| Imhof, W. L., Nakano, G. H. and Reagan, J. B.              | 1975b | J. Geophys. Res. <u>80</u> , 3629.  |
| Khan, J. M., Potter, D. L., and Worley, R. D.              | 1965  | Phys. Rev. <u>139</u> , A1735   |
| Luhmann, J. G. and Blake, J. B.                            | 1977  | J. Atmos. Terr. Phys. (in press).   |
| Madison, D. H. and Merzbacher, E.                          | 1975  | in Atomic Inner-Shell Processes, (ed. B. Crasemann) 1, Academic Press, New York.                                |
| Merzbacher, E. and Lewis, H. W.                            | 1958  | in Handbuch der Physik, (ed. S. Flügge), <u>34</u> , 66, Springer-Verlag, Berlin.                               |
| Tawara, H. Harrison, K. G., and de Heer, F. J.             | 1973  | Physica <u>63</u> , 351.  |



## THE IVAN A. GETTING LABORATORIES

The Laboratory Operations of The Aerospace Corporation is conducting experimental and theoretical investigations necessary for the evaluation and application of scientific advances to new military concepts and systems. Versatility and flexibility have been developed to a high degree by the laboratory personnel in dealing with the many problems encountered in the nation's rapidly developing space and missile systems. Expertise in the latest scientific developments is vital to the accomplishment of tasks related to these problems. The laboratories that contribute to this research are:

Aerophysics Laboratory: Launch and reentry aerodynamics, heat transfer, reentry physics, chemical kinetics, structural mechanics, flight dynamics, atmospheric pollution, and high-power gas lasers.

Chemistry and Physics Laboratory: Atmospheric reactions and atmospheric optics, chemical reactions in polluted atmospheres, chemical reactions of excited species in rocket plumes, chemical thermodynamics, plasma and laser-induced reactions, laser chemistry, propulsion chemistry, space vacuum and radiation effects on materials, lubrication and surface phenomena, photosensitive materials and sensors, high precision laser ranging, and the application of physics and chemistry to problems of law enforcement and biomedicine.

Electronics Research Laboratory: Electromagnetic theory, devices, and propagation phenomena, including plasma electromagnetics; quantum electronics, lasers, and electro-optics; communication sciences, applied electronics, semiconducting, superconducting, and crystal device physics, optical and acoustical imaging; atmospheric pollution; millimeter wave and far-infrared technology.

Materials Sciences Laboratory: Development of new materials; metal matrix composites and new forms of carbon; test and evaluation of graphite and ceramics in reentry; spacecraft materials and electronic components in nuclear weapons environment; application of fracture mechanics to stress corrosion and fatigue-induced fractures in structural metals.

Space Sciences Laboratory: Atmospheric and ionospheric physics, radiation from the atmosphere, density and composition of the atmosphere, aurorae and airglow; magnetospheric physics, cosmic rays, generation and propagation of plasma waves in the magnetosphere; solar physics, studies of solar magnetic fields; space astronomy, x-ray astronomy; the effects of nuclear explosions, magnetic storms, and solar activity on the earth's atmosphere, ionosphere, and magnetosphere; the effects of optical, electromagnetic, and particulate radiations in space on space systems.

THE AEROSPACE CORPORATION  
El Segundo, California

...

